#### **MONTEBAN** - narasin granule

Elanco Animal Health Co

**ELANCO** \*

AF0510-50B

For Use in Broiler

Chicken Feeds Only

Monteban<sup>®</sup> 45

Narasin

Net Weight: **50 lbs** (22.68 kg) **Type A Medicated Article** 

Do Not Feed Undiluted.

## **ACTIVE DRUG INGREDIENT**

For the prevention of coccidiosis caused by Eimeria necatrix, E. tenella, E. acervulina, E. brunetti, E. mivati, and E. maxima.

Important: Must be thoroughly mixed in feeds before use.

Mixing Directions – Thoroughly mix the following amounts of Monteban to provide 54 through 72 grams of narasin in one ton of feed. The dosage should be adjusted to meet the severity of the coccidial challenge, which varies with environmental and management conditions

It is recommended that a preblend of Monteban in 15-20 lbs of feed be made before incorporating into the total amount of finished feed.

Narasin	Monteban
(Grams/Ton)	(Lbs/Ton of Type C Feed)
	Lbs
54	1.20
59	1.31
63	1.40
68	1.51
72	1.60

**Feeding Directions**– Feed continuously as the sole ration.

**CAUTION:** Do not allow adult turkeys, horses or other equines access to narasin formulations. Ingestion of narasin by these species has been fatal.

NOT FOR HUMAN USE

WARNING: When mixing and handling Monteban, use protective clothing, impervious gloves and a dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.

Store in a cool, dry place.

Not to be used after the date printed at top of bag.



\*Elanco<sup>®</sup>, Monteban<sup>®</sup> and the diagonal color bar are trademarks of Eli Lilly and Company.

TAKE TIME



## **OBSERVE LABEL DIRECTIONS**

Restricted Drug (California) - Use Only a s Directed - NADA # 118-980, Approved by FDA

Elanco Animal Health

A Division of Eli Lilly and Company

Indianapolis, IN 46285, U.S.A.

To report adverse effects, access medical information, or obtain additional product information, call 1-800-428-4441. BG7070DEAMB (V01-02-2007)



Net Weight: 50 lbs

### Type A Medicated Article

# Do Not Feed Undiluted.

For the prevention of coccidiosis caused by Eimeria necatrix, E. tenella, E. acervulina, E. brunetti, E. mivati, and E. maxima. Important: Must be thoroughly mixed in feeds before use.

Mixing Directions—Thoroughly mix the following amounts of Monteban to provide 54 through 72 grams of narasin in one ton of feed. The dosage should be adjusted to meet the severity of the coccidial challenge, which varies with environmental and management conditions.

It is recommended that a preblend of Monteban in 15-20 lbs of feed be made before incorporating into the total amount of finished feed.


Narasin (Grams/Ton)	Monteban (Lbs/Ton of Type C Feed)
	Lbs
54	1.20
59	1.31
63	1.40
68	1.51
72	1.60

Feeding Directions-Feed continuously as the sole ration.

**CAUTION:** Do not allow adult turkeys, horses or other equines access to narasin formulations. Ingestion of narasin by these species has been fatal.

#### **NOT FOR HUMAN USE**

**WARNING:** When mixing and handling Monteban, use protective clothing, impervious gloves and a dust mask. Operators should wash thoroughly with soap and water after handling. If accidental eye contact occurs, immediately rinse thoroughly with water.





OBSERVE LABEL

\*Elanco®, Monteban® and the diagonal color bar are trademarks of Eli Lilly and Company.

Restricted Drug (California) - Use Only as Directed - NADA # 118-980, Approved by FDA

Elanco Animal Health A Division of Eli Lilly and Company Indianapolis, IN 46285, U.S.A.

To report adverse effects, access medical information, or obtain additional product information, call 1-800-428-4441.

BG7070DEAMB (V01-02-2007)



BG7070DEAMB